

NOTES

IS186 Insertion at a Hot Spot in the *lon* Promoter as a Basis for Lon Protease Deficiency of *Escherichia coli* B: Identification of a Consensus Target Sequence for IS186 Transposition

L. SAI SREE,¹ MANJULA REDDY,¹ AND J. GOWRISHANKAR^{1,2*}

Centre for Cellular and Molecular Biology, Hyderabad 500007,¹ and Centre for DNA Fingerprinting and Diagnostics, Hyderabad 500076,² India

Received 2 July 2001/Accepted 5 September 2001

The radiation sensitivity of *Escherichia coli* B was first described more than 50 years ago, and the genetic locus responsible for the trait was subsequently identified as *lon* (encoding Lon protease). We now show that both *E. coli* B and the first reported *E. coli* K-12 *lon* mutant, AB1899, carry IS186 insertions in opposite orientations at a single site in the *lon* promoter region and that this site represents a natural hot spot for transposition of the insertion sequence (IS) element. Our analysis of deposited sequence data for a number of other IS186 insertion sites permitted the deductions that (i) the consensus target site sequence for IS186 transposition is 5'-(G)_{≥4}(N)_{3–6}(C)_{≥4}-3', (ii) the associated host sequence duplication varies within the range of 6 to 12 bp and encompasses the N_(3–6) sequence, and (iii) in a majority of instances, at least one end of the duplication is at the G-N (or N-C) junction. IS186-related sequences were absent in closely related bacterium *Salmonella enterica* serovar Typhimurium, indicating that this IS element is a recent acquisition in the evolutionary history of *E. coli*.

The heritability of radiation sensitivity and radiation resistance traits in bacteria was first studied by Evelyn Witkin more than 50 years ago (26, 27), at a time when the genetic basis of inheritance in these organisms was still a matter of debate (14, 26, 27). Her choice of *Escherichia coli* B as the strain for these studies was historically significant, not least because it is the most radiation sensitive of all natural isolates of *E. coli* (14). In 1964, Howard-Flanders et al. (12) reported isolation of the first *E. coli* K-12 *lon* mutant, AB1899, and stated that “as regards sensitivity to radiation, . . . K-12 *lon*[−] resemble strains B. . .” That *E. coli* B (as well as its radiation-resistant suppressor derivative *E. coli* B/r [26, 27]) is also a natural *lon* mutant was shown subsequently by Donch and Greenberg (9). Deficiency of the Lon protease (encoded by *lon*) in the B strains is one reason that they have been developed as *E. coli* hosts of choice for overproduction of recombinant heterologous proteins (2, 24, 28).

We now report that the *lon* mutations in both *E. coli* B and AB1899 (*E. coli* K-12) were caused by IS186 insertions in opposite orientations at a single site within the spacer region of the *lon* promoter and that this site is a natural hot spot for IS186 transposition. An unusual feature of this insertion sequence (IS) element is that, unlike most other transposons (7, 25), it is associated with a variable length of duplication of the flanking host sequence at the sites of its insertion (5, 23).

Independent *lon*::IS186 insertions at a single site in four strains. We showed earlier (21) that K-12 strain GJ1823 had suffered a spontaneous and unselected IS186 insertion at the same site in the spacer region of the *lon* promoter as that identified by Ignatov and Chistyakova (13) on a plasmid derivative (pBLI) with the cloned *lon* gene. The genotype description for strain AB1899 at the *E. coli* Genetic Stock Center website (<http://cgsc.biology.yale.edu>) indicates, as a personal communication from D. A. Vlazny and C. W. Hill, that the *lon-1* mutation in the strain is caused by IS186 insertion although its position is not specified.

These facts provided us the rationale to compare, at the molecular level, the features of the *lon* mutation in strains AB1899, B/r, and GJ1823 with that published earlier for the plasmid pBLI by Ignatov and Chistyakova (13). We initially performed PCRs using pairwise combinations of the following three primers, designated A, B, and C, respectively: 5'-TGAC CAAGCAGTATCAGG-3', 5'-AAGATCGTTTACACCCGG CT-3', and 5'-GAGAAGCTGATTATATCGT-3'. Primers A and B were designed to amplify in PCR the region from −392 to +277 of *lon* (relative to the start site of *lon* transcription, taken as +1). Primer C is complementary to a site within the 1.34-kb IS186 element, situated 0.6 kb away from one of its ends. With primer pair A and B, the PCR product obtained with genomic DNA of wild-type *E. coli* strain MG1655 as the template was 1.3 kb smaller than that obtained with genomic DNA from each of the *lon* mutants AB1899, GJ1823, and B/r (Fig. 1). With the DNA templates from the three *lon* mutant strains, primer pair A and C on the one hand and pair B and C on the other behaved mutually exclusively, in that a PCR amplification product was obtained only with the latter for

* Corresponding author. Mailing address: Centre for DNA Fingerprinting and Diagnostics, ECIL Rd., Nacharam, Hyderabad 500076, India. Phone: 91-40-7155609. Fax: 91-40-7155610. E-mail: shankar@www.cdfd.org.in.




FIG. 3. Southern blot hybridization of genomic-DNA preparations from various strains using the *IS186* probe. Experimental methods were as described previously (22). The DNA preparations were digested with *EcoRV*, *BglI*, or *PvuII* as indicated and subjected to Southern blot hybridization with a ^{32}P -labeled *PstI*-*Bam*HI internal fragment of *IS186* (obtained from plasmid pHYD138 [21]), followed by autoradiography. Strains employed: 1, MG1655 (K-12 wild type); 2, B/r; 3, AB1899 (K-12 *lon*); 4, GJ1823 (K-12 *lon*); 5, BL21 (*E. coli* B); 6, *S. enterica* LT2. At the left are shown the positions of migration of DNA markers (sizes in kilobases). Arrows, *lon*-specific bands in the lanes corresponding to AB1899 and GJ1823; their sizes (5.6 [*EcoRV*] and 8.4 kb [*BglI*]) are consistent with those expected from *IS186* insertion in the cognate *lon*-bearing restriction fragments of wild-type K-12 (4, 20).

K-12 *lon*::*IS186* derivatives AB1899 and GJ1823 each have an additional copy which correlates with the insertion in *lon*. The two *E. coli* B derivatives each have five copies of *IS186*, the most plausible explanation for which is that they represent insertions in the loci homologous to the three of wild-type K-12, with additional copies in *lon* and near *hokX* (16). In another study, Birkenbihl and Vielmetter (3) also concluded that an *IS186* insertion into a new locus in a K-12 strain represented an additional (fourth) copy of the IS element.

In the Southern blot hybridization experiment (Fig. 3), we also included bacterial genomic DNA from strain LT2 of closely related organism *Salmonella enterica* serovar Typhimurium and observed no signal for the *IS186* probe. Our finding is consistent with the fact that *IS186* sequences have not so far been detected in the unfinished genome sequence of *S. enterica* (<http://www.tigr.org>). *IS186* has also not been detected in the genome of *E. coli* O157:H57 (10, 17) or in any of the other sequenced bacterial genomes, indicating that this IS element, much like others such as IS2, IS4, and IS30 (8), may be a recent acquisition in the evolutionary history of *E. coli*. The findings of Pedersen and Gerdes (16) suggest that *IS186* is present in *E. coli* C and also in not less than 44 of the 72 natural *E. coli* isolates of the ECOR collection.

Consensus sequence for *IS186* transposition. Different IS elements are known to vary in the extent to which they exhibit target site selectivity for transposition (7, 25). A consensus sequence for *IS186* transposition has not previously been determined, although it is known that the insertions occur in G-C rich target sequences (5, 15, 23, 25). Because many eukaryotic sequences cloned in *E. coli* hosts are G-C rich, several instances of *IS186* insertions in them are identified in the GenBank database. We analyzed all the deposited entries to deduce that sequence 5'-(G) \geq 4(N) \geq 3(C) \geq 4-3' is the consensus target recognition site for *IS186* transposition (Table 1). The

lon hot spot fits the consensus, as do all but one of the entries listed in Table 1. Our analysis also indicates that, even as its length varies between 6 and 12 bp, the associated host sequence duplication always encompasses the (N) \geq 3 sequence of the target recognition site. Furthermore, in more than half of the instances examined at least one end of the duplication (that is, at least one site of endonucleolytic cleavage during transposition) is at the G-N (or N-C) junction (Table 1).

These findings may offer approaches to study the mechanism of *IS186* transposition, of which at present little is known. Our analysis suggests that, in *IS186* transposition, there is an initial specificity in target sequence recognition by transposase but that the sites where subsequent endonucleolytic cleavages occur are less constrained; such a phenomenon invites comparisons with both the mechanism of type I restriction endonuclease action (18) and that of rearrangements of the immunoglobulin genes mediated by the RAG1 and RAG2 proteins, where a preference for G-C-rich sequences at the target sites (11) and the generation of duplications of variable length (1) have also been observed.

It has been suggested that *IS186* belongs to the *IS4* family of insertion elements (5, 15, 19, 25), but the family as currently defined includes members that generate variable (*IS4* and *IS186*) as well as fixed lengths of target site duplications. The length of the duplications associated with transposition of *IS50* (Tn5), which is another member of the *IS4* family, is usually 9 bp but rarely could also be 10 bp (6). Finally, *IS1* is also known to generate host sequence duplications of various lengths (25) but is not a member of the *IS4* family.

We thank Mary Berlyn for bacterial strains and acknowledge the assistance of Mehar Sultana and N. Nagesh with oligonucleotide synthesis and DNA sequencing, respectively. We also thank one of the anonymous reviewers for suggesting the analogy between *IS186* transposition on the one hand and restriction endonuclease action and antibody gene rearrangement on the other.

J. G. is an honorary faculty member of the Jawaharlal Nehru Centre for Advanced Scientific Research.

REFERENCES

- Agrawal, A., Q. M. Eastman, and D. G. Schatz. 1998. Transposition mediated by RAG1 and RAG2 and its implications for the evolution of the immune system. *Nature* **394**:744-751.
- Bhandari, P., and J. Gowrishankar. 1997. An *Escherichia coli* host strain useful for efficient overproduction of cloned gene products with NaCl as the inducer. *J. Bacteriol.* **179**:4403-4406.
- Birkenbihl, R. P., and W. Vielmetter. 1991. Completion of the IS map in *E. coli*: *IS186* positions on the *E. coli* K12 chromosome. *Mol. Gen. Genet.* **226**:318-320.
- Blattner, F. R., G. Plunkett III, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453-1462.
- Chong, P., I. Hui, T. Loo, and S. Gillam. 1985. Structural analysis of a new GC-specific insertion element *IS186*. *FEBS Lett.* **192**:47-52.
- Chu, C. C., and A. J. Clark. 1989. A 10- rather than 9-bp duplication associated with insertion of Tn5 in *Escherichia coli* K-12. *Plasmid* **22**:260-264.
- Craig, N. L. 1996. Transposition, p. 2339-2362. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. American Society for Microbiology, Washington, D.C.
- Deonier, R. C. 1996. Native insertion sequence elements: locations, distributions, and sequence relationships, p. 2000-2011. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. American Society for Microbiology, Washington, D.C.

9. Donch, J., and J. Greenberg. 1968. Ultraviolet sensitivity gene of *Escherichia coli* B. *J. Bacteriol.* **95**:1555–1559.
10. Hayashi, T., K. Makino, M. Ohnishi, K. Kurokawa, K. Ishii, K. Yokoyama, C.-G. Han, E. Ohtsubo, K. Nakayama, T. Murata, M. Tanaka, T. Tobe, T. Iida, H. Takami, T. Honda, C. Sasakawa, N. Ogasawara, T. Yasunaga, S. Kuhara, T. Shiba, M. Hattori, and H. Shinagawa. 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* **8**(Suppl.):47–52.
11. Hiom, K., M. Melek, and M. Gellert. 1998. DNA transposition by the RAG1 and RAG2 proteins: a possible source of oncogenic translocations. *Cell* **94**:463–470.
12. Howard-Flanders, P., E. Simson, and L. Theriot. 1964. A locus that controls filament formation and sensitivity to radiation in *Escherichia coli* K-12. *Genetics* **49**:237–246.
13. Ignatov, K. B., and L. G. Chistyakova. 1997. Integration of the insertion element IS186 into the –10 region of the heat shock promoter of the *Escherichia coli lon* gene. *Bioorg. Khim.* **23**:18–20. (In Russian.)
14. Kelner, A., W. D. Bellamy, G. E. Stapleton, and M. R. Zelle. 1955. Symposium on radiation effects on cells and bacteria. *Bacteriol. Rev.* **19**:22–44.
15. Kothary, R. K., D. Jones, and E. P. M. Candido. 1985. IS186: an *Escherichia coli* insertion element isolated from a cDNA library. *J. Bacteriol.* **164**:957–959.
16. Pedersen, K., and K. Gerdes. 1999. Multiple *hok* genes on the chromosome of *Escherichia coli*. *Mol. Microbiol.* **32**:1090–1102.
17. Perna, N. T., G. Plunkett III, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grotbeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamouisis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner. 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**:529–533.
18. Redaschi, N., and T. A. Bickle. 1996. DNA restriction and modification systems, p. 773–781. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. American Society for Microbiology, Washington, D.C.
19. Rezsöhazy, R., B. Hallet, J. Delcour, and J. Mahillon. 1993. The IS4 family of insertion sequences: evidence for a conserved transposase motif. *Mol. Microbiol.* **9**:1283–1295.
20. Rudd, K. E. 1998. Linkage map of *Escherichia coli* K-12, edition 10: the physical map. *Microbiol. Mol. Biol. Rev.* **62**:985–1019.
21. SaiSree, L., M. Reddy, and J. Gowrishankar. 2000. *lon* incompatibility associated with mutations causing SOS induction: null *uvrD* alleles induce an SOS response in *Escherichia coli*. *J. Bacteriol.* **182**:3151–3157.
22. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
23. Sengstag, C., S. Iida, R. Hiestand-Nauer, and W. Arber. 1986. Terminal inverted repeats of prokaryotic transposable element IS186 which can generate duplications of variable length at an identical target sequence. *Gene* **49**:153–156.
24. Studier, F. W., and B. A. Moffatt. 1986. Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *J. Mol. Biol.* **189**:113–130.
25. Syvanen, M. 1988. Bacterial insertion sequences, p. 331–356. In R. Kucherlapati and G. R. Smith (ed.), *Genetic recombination*. American Society for Microbiology, Washington, D.C.
26. Witkin, E. M. 1946. Inherited differences in sensitivity to radiation in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **32**:59–68.
27. Witkin, E. M. 1947. Genetics of resistance to radiation in *Escherichia coli*. *Genetics* **32**:221–248.
28. Young, R. A., and R. W. Davis. 1983. Efficient isolation of genes by using antibody probes. *Proc. Natl. Acad. Sci. USA* **80**:1194–1198.