

1. Banerjee, P. K., Geology and geochemistry of the Sukinda ultramafic field, Cuttack district, Orissa. *Mem. Geol. Surv. India*, 1972, **103**, 17.
2. Chakraborty, K. L. and Chakraborty, T. L., Geological features and origin of the chromite deposits of Sukinda valley, Orissa, India. *Miner. Deposita*, 1984, **19**, 256–265.
3. Page, N. J., Banerjee, P. K. and Haffty, J., Characterization of the Sukinda and Nausahi ultramafic complex, Orissa, India by platinum-group element geochemistry. *Precambrian Res.*, 1985, **30**, 27–41.
4. Varma, O. P., Some aspects of ultramafic and ultrabasic rocks and related chromite metallogenesis with examples from eastern region of India. In Proceedings of the Seventy-third Session of the Indian Science Congress Association, Delhi, 1986, pp. 1–72.
5. Mondal, S. K., Ripley, E. M., Li, C. and Frei, R., The genesis of Archaean chromitites from the Nuasahi and Sukinda massifs in the Singhbhum Craton, India. *Precambrian Res.*, 2006, **148**, 45–66.
6. Vermaak, C. F., A brief overview of South Africa's mineral industry: World context and changing local circumstances. *Miner. Deposita*, 1997, **32**, 312–322.
7. Sarkar, N. K., Panigrahi, D., Ghosh, S. N. and Mallik, A. K., A note on the incidence of gold-PGM in the breccia zone of Katpal chromium quarry, Sukinda ultramafic complex, Dhenkal Dt., Orissa *Indian Miner.*, 2003, **57**, 85–92.
8. Sen, A. K., Sharma, P. K., Mohanthy, D. and Ghosh, T. K., Composition of Cr spinel – An ore genetic indicator of Katpal chromite deposit, Sukinda ultramafic field, Orissa. *Curr. Sci.*, 2005, **88**, 1547–1550.
9. Basu, A., Maitra, M. and Roy, P. K., Petrology of mafic-ultramafic complex of Sukinda valley, Orissa, *Indian Miner.*, 1997, **50**, 271–290.
10. Auge, T. and Lerouge, C., Mineral-chemistry and stable-isotope constraints on the magmatism, hydrothermal alteration, and related PGE – (base-metal sulphide) mineralisation of the Mesoproterozoic Baula-Nuasahi Complex. *Miner. Deposita*, 2004, **39**, 583–607.
11. Sarkar, N. K., Mallik, A. K., Panigrahi, D. and Ghosh, S. N., A note on the occurrence of breccia zone in the Katpal chromite lode, Dhenkanal district, Orissa. *Indian Miner.*, 2001, **55**, 247–250.
12. Burkhard, D. J. M., Accessory chromium spinels: Their coexistence and alteration in serpentinites. *Geochim. Cosmochim. Acta*, 1993, **57**, 1297–1306.
13. Mohanthy, J. K., Sahoo, R. K. and Paul, A. K., Chromite alteration at Baula-Nuasahi igneous complex, Orissa. *J. Geol. Soc. India*, 1996, **48**, 265–276.
14. Naldrett, A. J. and Cabri, L. J., Ultramafic and related mafic rocks: Their classification and genesis with special reference to the concentration of nickel sulphides and platinum-group elements. *Econ. Geol.*, 1976, **71**, 1131–1158.
15. Arndt, N. T., Leshner, C. M. and Czamanske, G. K., Mantle-derived magmas and magmatic Ni–Cu–(PGE) deposits. *Econ. Geol.*, 2005, 100th Anniversary Volume, 5–23.
16. Naldrett, A. J. and Lehmann, J., Spinel non-stoichiometry as the explanation for Ni-, Cu-, and PGE-enriched sulphides in chromitites. In *Geo Platinum '87*, Elsevier, London, pp. 93–110.
17. Merkle, R. K. W., Platinum-group minerals in the middle group of chromitite layers at Marikana, western Bushveld Complex – Indications for collection mechanisms and postmagmatic modification. *Can. J. Earth Sci.*, 1992, **29**, 209–221.
18. Crockett, J. H., Platinum-group element geochemistry of mafic and ultramafic rocks. In *The Geology, Geochemistry, Mineralogy and Mineral Beneficiation of Platinum Group Elements* (ed. Cabri, L. J.), Canadian Institute of Mining, Metallurgy and Petroleum, 2002, vol. 54, pp. 117–210.
19. Von Gruenewaldt, G. and Merkle, R. K. W., Platinum group element proportions in chromitites of the Bushveld complex: Implications for fractionation and magma mixing models. *J. Afr. Earth Sci.*, 1995, **21**, 615–632.
20. Rajamani, V. and Naldrett, A. J., Partitioning of Fe, Co, Ni, and Cu between sulphide liquid and basaltic melts and the composition of Ni–Cu sulphides. *Econ. Geol.*, 1978, **73**, 82–93.
21. Makovicky, M., Makovicky, E. and Rose-Hansen, J., Experimental studies on the solubility and distribution of platinum group elements in base-metal sulphides in platinum deposits. In *Metallogeny of Basic and Ultrabasic Rocks* (eds Gallagher, M. J. et al.), The Institute of Mining and Metallurgy, London, 1986, pp. 415–425.
22. Fleet, M. E., Crockett, J. H. and Stone, W. E., Partitioning of platinum-group elements (Os, Ir, Ru, Pt, Pd) and gold between sulfide liquid and basalt melt. *Geochim. Cosmochim. Acta*, 1996, **60**, 2397–2412.

ACKNOWLEDGEMENTS. The Department of Science and Technology, New Delhi, is acknowledged for providing financial support to carry out this work. The Managing Director of OMC is acknowledged for permission to carry out this scientific study. The Director, NGRI is acknowledged for support and guidance. The work was carried out while P.V.S.R. was a BOYSCAST fellow at the Department of Geology, University of Pretoria, South Africa.

Received 1 December 2006; revised accepted 27 June 2007

A new nightfrog, *Nyctibatrachus minimus* sp. nov. (Anura: Nyctibatrachidae): The smallest frog from India

S. D. Biju^{1,2,*}, Ines Van Bocxlaer²,
Varad B. Giri³, Kim Roelants², J. Nagaraju⁴ and
Franky Bossuyt²

¹Centre for Environmental Management of Degraded Ecosystems, School of Environmental Studies, University of Delhi 110 007, India

²Biology Department, Unit of Ecology and Systematics, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium

³Herpetology Section, Bombay Natural History Society (BNHS), S.B. Singh Road, Mumbai 400 023, India

⁴Laboratory of Molecular Genetics, Centre for DNA Fingerprinting and Diagnostics, ECIL Road, Nacharam, Hyderabad 500 076, India

A new nightfrog, *Nyctibatrachus minimus* sp. nov. (Anura: Nyctibatrachidae) is described from Kurichiyarmala in the Western Ghats, India. Its most distinctive feature is the small adult snout-vent length, averaging only 12.3 mm in adult males ($N = 15$), making it the smallest known frog from India. Analyses of a fragment of the mitochondrial NADH dehydrogenase 1 gene indicate a minimum divergence of 22% with known small-sized congeners. Miniaturization in *Nyctibatrachus* sp. seems to be associated with absence of

*For correspondence. (e-mail: sdbiju@cemde.du.ac.in)

webbing on toes and fingers, which may have resulted from evolutionary specialization to life in terrestrial habitats.

Keywords: NADH dehydrogenase 1, *Nyctibatrachus minimus* sp. nov., smallest Indian frog, Western Ghats.

THE Western Ghats of Peninsular India is considered an important amphibian hotspot, both in terms of species richness and higher-taxonomic-level endemism^{1–5}. The genus *Nyctibatrachus* Boulenger, 1882 is an ancient, endemic lineage⁵ in the family Nyctibatrachidae⁶ that currently holds 12 recognized species^{7,8}. We herein describe a miniature species from Kurichiyarmala, *Nyctibatrachus minimus* sp. nov. and compare it with the descriptions and available material of all species currently recognized in the genus.

N. minimus sp. nov. Holotype: Bombay Natural History Society (BNHS), Mumbai, Maharashtra, India, BNHS 4527, an adult male collected by S.D.B. on 28 July 2001 at an altitude of 1200 m asl, from Kurichiyarmala, 11°35'N, 75°58'E, Wayanad District, Kerala, India; Paratypes: BNHS 4528–4534 (seven adult males), collected along with the holotype, and BNHS 4535–4540 (six adult males) and BNHS 4541 (an adult female), collected by S.D.B. on 24 February 2006 from the type locality. The species epithet *minimus* (Latin: smallest) refers to the diminutive adult size. Suggested vernacular name: miniature nightfrog.

N. minimus sp. nov. can be distinguished from all other members of the genus by the combination of small adult size (males: range 10.0–14.0 mm, $N = 15$; female: 14.9 mm, $N = 1$), presence of granular femoral glands in breeding males, presence of two rows of weakly developed vomerine teeth on each side, lack of webbing on hands and feet, and relatively smooth dorsal skin with faint and interrupted dorsolateral folds and glandular projections (Figures 1–4).

Comparisons were made with the available type specimens of each name bearing taxon in *Nyctibatrachus*, and with other recent collections from the type localities made by the authors. Sexual maturity was determined by examining gonads through a small lateral incision, and by the presence of an advertisement call and femoral glands in breeding males. Nine *Nyctibatrachus* species (*N. aliciae*, *N. deccanensis*, *N. humayuni*, *N. karnatakaensis*, *N. major*, *N. petraeus*, *N. sanctipalustris*, *N. sylvaticus*, and *N. vasanthi*) are clearly distinct from *N. minimus* sp. nov. by their larger adult snout-vent length and relatively robust body. *Nyctibatrachus minimus* sp. nov. can be distinguished from the small-sized *N. kempholeyensis*, a species that has not been observed since its original description⁹, by its smaller snout-vent size (maximum 14.9 mm in female *N. minimus* vs 18.0 mm in original description of *N. kempholeyensis*), head about as long as broad (vs head wider than long), and loreal region acute

to slightly vertical (vs loreal region vertical to oblique). To support the distinction of *N. minimus* sp. nov. from the remaining smaller *Nyctibatrachus* species, we sequenced a fragment of ~625 base pairs of the mitochondrial NADH dehydrogenase 1 (*NDI*) gene. DNA sequences were obtained by whole-genome extraction¹⁰, PCR amplification, and cycle-sequencing along both strands (GenBank accession numbers EF136395–EF136398 for BNHS 4540, *N. minimus* sp. nov.; BNHS 4520, *N. anamallaiensis*; SDB 1149, *N. beddomii*; BNHS 4522, *N. minor* respectively). Uncorrected pairwise comparison of the sequence of *N. minimus* sp. nov. with those of *N. beddomii*, *N. anamallaiensis*¹¹ (currently considered a synonym of *N. beddomii*^{12–14}), and *N. minor* shows genetic divergences of 25%, 27% and 22% respectively. A 22% sequence divergence between *N. beddomii* and *N. anamallaiensis*, together with a dissimilar morphology (the most conspicuous difference being the presence of dorsolateral folds in *N. anamallaiensis*), allow us to remove the latter from synonymy of *N. beddomii*. Morphologically, *N. minimus* sp. nov. differs from *N. anamallaiensis*, *N. beddomii* and *N. minor* by its still smaller adult size, whose range shows




Figure 1. *Nyctibatrachus minimus* sp. nov. **a**, Holotype (BNHS 4527) in life. **b**, Holotype in life (ventral side), showing femoral glands. **c**, Paratype (BNHS 4528) in preservation (ventral side), showing femoral glands.

little or no overlap with those of the other species (Figure 4, Table 1), and by its dorsal skin having faint and interrupted dorsolateral folds and glandular projections (vs distinct dorsolateral folds separated by a row of glandular

projections in *N. anamallaiensis*, distinct dorsolateral glandular folds separated by an 'X' pattern on anterior half of back in *N. minor*, and a smooth dorsal skin in *N. beddomii*). In addition, *N. minimus* sp. nov. differs from *N. anamallaiensis* by its weakly developed vomerine teeth (vs well-developed vomerine teeth) and from *N. minor* by the presence of femoral glands (vs absence of femoral glands)¹².

The description (all measurements in mm) of the holotype follows terminology used elsewhere¹⁵. Small-sized frog (SVL 13.2); head (Figure 2) about as long as broad (HW 5.6; HL 5.5); outline of snout in dorsal view oval, its length (SL 1.8) larger than the horizontal diameter of the eye (EL 1.2); canthus rostralis indistinct, loreal region acute to slightly vertical; interorbital area (IUE 2.3) larger than upper eyelid (UEW 0.7); tympanum indistinct; vomerine teeth weakly developed, pointed, present in two rows on each side, oblique between choanae; tongue nearly cordate, emarginated, without a papilla; button-like projection at the base of the tongue on lower jaw; supratympanic fold absent. Forelimb length (FLL 2.3) slightly shorter than hand (HAL 2.9); tips of fingers with weakly developed disks, with marginal grooves; webbing on fingers absent; subarticular tubercle rather prominent, rounded, single, IV1 absent (Figure 3 a), longitudinal groove on the dorsal surface of finger disks; prepollex oval and rather indistinct; supernumerary tubercles absent. Hindlimbs moderately long; shank 3.3 times longer (ShL 6.6) than wide (ShW 2.0), almost equal to femur length (FL 6.5), longer than distance from base of inner metatarsal tubercle to tip of toe IV (FOL 5.9), 1.4 times




Figure 2. *Nyctibatrachus minimus* sp. nov. holotype. a, Dorsal view; b, Ventral view.




Figure 3. *Nyctibatrachus minimus* sp. nov.. a, Ventral view of hand; b, Ventral view of foot; c, Magnified dorsal view of toe showing groove.




Figure 4. Comparison of snout-vent length of males of the four smallest species of *Nyctibatrachus*. Measurements are summarized in box and whisker plots, with the box spanning the interquartile range, the median indicated by a line inside the box, and the whiskers extending to the highest and lowest observations. From left to right: *N. minimus* sp. nov. ($N = 15$), *N. anamallaiensis* ($N = 10$), *N. beddomii* ($N = 10$) and *N. minor* ($N = 10$).

Table 1. Morphometric measurements (all in mm; range, mean and standard deviation) of *Nyctibatrachus minimus* sp. nov., *N. anamallaiensis*, *N. beddomii* and *N. minor*. Abbreviations are defined in the text. Specimens used in analyses (BNHS; NHM, Natural History Museum, London; SDB, field number: housed at CEMDE, University of Delhi): *N. minimus* sp. nov. – males: BNHS 4527–4540, SDB 421 (all from Kurichiyarmala); female BNHS 4541 (Kurichiyarmala); *N. anamallaiensis* – males: BNHS 4256 and BNHS 4520 (Andiparai Shola), NHM 74.4.29.1372–1373 ('Anamallays'), SDB 40319, SDB 5611–5615 (Puthuthottam); female BNHS 4521 (Andiparai Shola); *N. beddomii* – males: NHM 74.4.29.519, NHM 74.4.29.457 (Malabar), NHM 82.2.10.31 ('Tivively'), BNHS 4252 (Ponmudi), BNHS 4253–4254 (Agasthyamala–Athirimala), BNHS 4255 (Ponmudi), SDB 4623 (Sengaltheri), SDB 1149–1150 (Ponmudi); female: NHM 74.4.29.456; *N. minor* – males: BNHS 4522–4526 (Ponmudi), SDB 401, SDB 1154–1155, SDB 6322, SDB 009 (all from Ponmudi); female: SDB 010 (Ponmudi)

	<i>N. minimus</i> sp. nov.				<i>N. anamallaiensis</i>				<i>N. beddomii</i>				<i>N. minor</i>			
	Male (N = 15)			Female N = 1	Male (N = 10)			Female N = 1	Male (N = 10)			Female N = 1	Male (N = 10)			Female N = 1
	Range	Mean	SD		Range	Mean	SD		Range	Mean	SD		Range	Mean	SD	
SVL	10.0–14.0	12.3	1.4	14.9	14.0–15.7	14.9	0.7	17.0	13.5–15.6	14.6	0.6	18.6	15.7–18.7	17.4	0.8	21.8
HW	4.2–5.8	5.2	0.5	5.9	5.1–5.7	5.5	0.2	6.4	5.1–6.5	5.6	0.5	7.1	6.7–8.1	7.3	0.4	10.0
HL	4.0–5.5	5.0	0.5	5.4	4.1–4.9	4.5	0.3	5.6	4.4–6.1	5.1	0.6	0.7	6.2–7.8	6.6	0.7	8.9
SL	1.7–2.5	2.0	0.2	2.4	2.0–2.1	2.0	0.1	2.3	2.1–2.5	2.3	0.1	2.6	2.8–3.5	3.2	0.2	4.3
EL	1.2–1.7	1.5	0.1	1.7	2.4–2.6	2.5	0.1	2.7	1.3–2.2	1.7	0.3	2.3	1.9–2.3	2.0	0.1	3.8
IUE	1.4–2.3	1.8	0.3	2.0	1.5–2.2	1.9	0.2	2.1	1.9–2.3	2.0	0.2	2.4	2.0–2.5	2.3	0.2	3.4
UEW	0.5–0.9	0.7	0.1	0.9	0.8–1.8	1.3	0.4	1.9	0.8–1.1	0.9	0.1	1.3	0.8–1.1	0.9	0.1	1.7
FLL	2.0–2.9	2.5	0.3	2.6	2.4–2.8	2.7	0.2	3.9	2.3–3.4	2.8	0.3	3.7	2.7–3.5	3.0	0.3	4.8
HAL	2.5–3.8	3.1	0.4	3.5	2.9–3.5	3.2	0.2	4.1	2.4–3.1	2.9	0.4	3.9	3.9–4.5	4.3	0.2	5.2
ShL	4.9–6.8	6.1	0.7	6.8	6.3–7.4	6.8	0.4	7.8	5.9–7.4	6.7	0.5	7.8	8.1–9.2	8.6	0.3	10.3
FL	4.2–6.6	5.8	0.7	5.9	5.9–7.6	6.8	0.5	7.8	5.4–7.4	6.5	0.6	7.6	8.1–9.1	8.7	0.3	10.5
TFOL	7.0–11.6	9.8	1.3	11.2	10–10.6	10.3	0.2	11.7	9.1–12.8	10.9	1.2	12.9	12.0–14.2	12.8	0.7	16.5
FOL	4.7–7.6	6.0	0.8	6.5	6.5–7.5	6.9	0.3	8.4	5.3–8.1	6.5	0.8	8.3	7.7–9.2	8.4	0.4	11.0




Figure 5. Distribution map for the four smallest species of *Nyctibatrachus* from the Western Ghats.

in distance from heel to tip of toe IV (TFOL 9.4); tips of toes with weakly developed disks, with marginal grooves; webbing absent (Figure 3 b); subarticular tubercles rather

distinct, rounded, simple, IV2, 3 and V1 absent (Figure 3 b); inner metatarsal tubercle indistinct, oval; supernumerary tubercles absent; longitudinal groove on the dorsal surface of toe disks (Figure 3 c). Skin of snout shagreened to granular, upper eyelids with a few prominent tubercles; side of head, anterior and posterior parts of dorsum and upper and lower parts of flank granular, with a chain of granular projections forming a faint and interrupted dorso-lateral fold from behind the eye to the vent on either side (Figures 1 a and 2 a); ventral side smooth. A pair of prominent oval-shaped femoral glands (length 3.1 mm × width 1.2 mm) is present on the hind legs (Figures 1 b and 2 b).

Colour of holotype in alcohol: A dark gray dorsum; dorsal surface of forelimbs and hindlimbs light grey with dark brown cross bands; ventrally uniform greyish with irregular spots, forearm and shank with brownish black markings.

Colour of holotype in life: A light tan dorsum without prominent patterns, except for a few dark brown markings between the eyes; iris light greyish-brown; forelimbs and hindlimbs light greyish-brown, with faint cross-bands; side of the head and lateral side with scattered minute white spots (Figure 1 a); ventrally light reddish-grey, forelimbs and hindlimbs with grey spots forming a vermiculated pattern; femoral glands orange (Figure 1 b).

Variation: Measurements (range, mean and standard deviation) of the type series are given in Table 1. Femoral glands of male specimens collected during the non-breeding season (February) were weakly developed compared to those of breeding males (July).

Distribution and natural history: *N. minimus* sp. nov. is currently known only from the type locality Kurichiyarmala, north of the Palghat Gap (Figure 5). This region is characterized by Shola forests (patches of high-altitude forest separated from one another by undulating grassland). Males start calling from marshes inside the forest during or immediately after sunset, with two subgular, external vocal sacs laterally inflated.

Size differentiation among nyctibatrachid species has been invoked to warrant recognition of the genus *Nanobatrachus* (*Nano* = small, *Batrachus* = frog)¹⁶. However, after the description of *N. minor*¹², a small species with typical *Nyctibatrachus* features, subsequent authors considered the two genera to be synonyms^{13,14}. With the present description of *N. minimus* sp. nov. and removal of *N. anamallaiensis* from the synonymy of *N. beddomii*, the genus *Nyctibatrachus* now contains 14 species. *N. minimus* sp. nov., *N. anamallaiensis* and *N. beddomii* are among the smallest of approximately 220 currently known Indian species of frogs⁶. The lack of webbing on their feet contrasts with the extensively webbed toes of larger species. Furthermore, while most species of *Nyctibatrachus* are aquatic, the miniature species mainly inhabit the forest floor, where they are mostly found in leaf litter or under rocks. Miniaturization in Nyctibatrachidae may thus have been associated with a morphological specialization to life in terrestrial habitats.

1. Inger, R. F., Distribution of amphibians in southern Asia and adjacent islands. In *Patterns of Distribution of Amphibians: A Global Perspective* (ed. Duellman, W. E.), The Johns Hopkins University Press, Baltimore, 1999, pp. 445–482.
2. Biju, S. D., A synopsis to the frog fauna of the Western Ghats, India. *Indian Soc. Conser. Biol. Publ.*, 2001, **1**, 1–24.
3. Biju, S. D. and Bossuyt, F., New frog family from India reveals an ancient biogeographical link with the Seychelles. *Nature*, 2003, **425**, 711–714.
4. Van Bocxlaer, I., Roelants, K., Biju, S. D., Nagaraju, J. and Bossuyt, F., Late Cretaceous vicariance in gondwanan amphibians. *PLoS ONE*, 2006, **1**, e74.
5. Roelants, K., Jiang, J. and Bossuyt, F., Endemic ranid (Amphibia: Anura) genera in southern mountain ranges of the Indian subcontinent represent ancient frog lineages: Evidence from molecular data. *Mol. Phylogenet. Evol.*, 2004, **31**, 730–740.

6. Frost, D. R., Amphibian species of the world: An online reference, version 5.0. American Museum of Natural History, New York, USA, 1 February 2007. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>
7. Das, I. and Kunte, K., New species of *Nyctibatrachus* (Anura: Ranidae) from Castle Rock, Karnataka State, Southwest India. *J. Herpetol.*, 2005, **39**, 465–470.
8. Dinesh, K. P., Radhakrishnan, C., Manjunatha Reddy, A. H. and Gururaja, K. V., *Nyctibatrachus karnatakaensis* nom. nov., a replacement name for the giant wrinkled frog from the Western Ghats. *Curr. Sci.*, 2007, **93**, 246–250.
9. Rao, C. R. N., On some new forms of Batrachia from S. India. *Proc. Indian Acad. Sci.*, 1937, **6**, 387–427.
10. Sambrook, J., Fritsch E. F. and Maniatis, T., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York, 1989.
11. Myers, E., A new frog from the Anamalai Hills, with notes on other frogs and some snakes from south India. *Proc. Biol. Soc. Wash.*, 1942, **55**, 49–56.
12. Inger, R. F., Shaffer, H. B., Koshy, M. and Bakde, R., A report on a collection of amphibians and reptiles from the Ponmudi, Kerala, South India. *J. Bombay Nat. Hist. Soc.*, 1984, **81**, 406–570.
13. Dubois, A., *Miscellanea taxonomica batrachologica I. Alytes*, 1986, **5**, 7–59.
14. Shaffer, H. B., Size and scaling in the Indian frogs *Nyctibatrachus* and *Nanobatrachus* (Ranidae). *Fieldiana, Zool. New Ser.*, 1988, **46**, 1–10.
15. Biju, S. D. and Bossuyt, F., A new species of frog (Ranidae, Rhacophorinae, *Philautus*) from the rainforest canopy in the Western Ghats, India. *Curr. Sci.*, 2005, **88**, 176–180.
16. Boulenger, G. A., *Catalogue of the Batrachia Salientia s. Ecaudata in the Collection of the British Museum*, London, 1882.

ACKNOWLEDGMENT. We are grateful to M. Wilkinson and B. T. Clarke (NHM, London) for providing access to specimens in their care, and D. J. Gower (NHM) and D. San Mauro (MNCN) for helpful comments on an earlier version of the manuscript. S.D.B. is grateful to the Indian National Science Academy, New Delhi and The Royal Society of London for a fellowship (2003) for travel and museum studies; the Kerala Forest Department for collection permits (No. WL-11-1558/01), and Anil Zachariya and K. Jayaram for assistance with the field studies. I.V.B. is supported by the Institute for the Promotion of Innovation through Science and Technology in Flanders (IWT-Vlaanderen). F.B. and K.R. are postdoctoral researchers at the Fonds voor Wetenschappelijk Onderzoek (FWO-Vlaanderen).

Received 29 May 2006; revised accepted 28 June 2007